

Exploring The Level of Awareness Among the Parents of Beta Thalassemia Patients in District Swat

Mehnaz bibi¹, Amir Sultan², Zaifullah³

Abstract: Background: Thalassemia is an inherited condition that affects globin chain synthesis and affects hemoglobin production in various ways. The aim of the study was to determine the level of awareness of parents of patients diagnosed with beta-thalassemia (βT).

Methodology: A cross-sectional descriptive study design was used to collect data from parents of patient regularly visiting blood transfusion centers in Swat district from February to March 2024. The sample size of the study was 235 using purposive sampling technique, with data collected through a reliable and valid questionnaire. The study was approved by the Ethics Review Board, with informed consent obtained from each participant. Data were analyzed using SPSS 22 as descriptive and inferential statistics.

Results: In the current study, most of the parents were female 146 (62.1%), age group 31-40 years 164 (69.8%), house wife 67 (28.5%), with a monthly income of 21,000 to 50,000 107 (45.5%), and parents without education with a civil registry 60 (25.5%). For most patients, the level of awareness was good (42%), followed by poor awareness (31%), while 27% of participants' responses were neutral. Most parents are aware of prohibited foods, thalassemia symptoms, preventive tests, complications, drug avoidance, regular laboratory tests, thalassemia genetic disorder, blood related marriages and HIV as a blood borne disease.

Conclusion: The study concluded that majority of the participant level of awareness was good, but a national basis campaign is required that address the preventive strategies to prevent thalassemia.

Keywords: Thalassemia, awareness, parents, Prevention, Transmission, Blood Transfusion.

1. BSN Student, Tasleem college of Nursing and health sciences swat, KPK
2. Assistant professor/ Principal: Tasleem college of Nursing and health sciences, swat KPK
3. MSN Scholar, School of nursing and midwifery, Aga khan university Karachi

Corresponding Author: Amir Sultan,
Assistant Professor / Principal:
Tasleem college of Nursing and health
sciences, swat KPK,
Email: amirsultan204@gmail.com

INTRODUCTION:

Thalassemia are inherited blood illnesses that minimize the level of hemoglobin (Hb) production, that are divided as alpha (α) and beta (βT). It is common among Italian, Greek, Middle Eastern, South Asian, and African people (Thalassemias. 2012). Alpha and βT are frequently passed down in an inherited manner. A child cannot develop thalassemia unless both parents are carriers. Parents with trait carrier of Thalassemia can result as twenty five percent of possibility may affect the baby in pregnancy that leads to hemoglobinopathy.

Thalassemia was reported in 280 million people worldwide in 2015, accounting for 16,800 fatalities (Hay et al. 2017) (Uthman, 2016). βT is a prevalent single-gene hereditary disease worldwide (Cousens et al. 2010). Every year, around 70,000 newborns worldwide are born with βT , and 270 million people carry haemoglobinopathies, according to various research. Thalassaemia is a public health issue worldwide because of its high frequency (Ishaq et al. 2012). βT is a hereditary condition that affects globin chain synthesis in varying ways. This disease is avoidable, as proved in Italy,

Greece, and Cyprus (Ansari et al. 2012).

Pakistan is among the low-income countries with high population and with no reliable data on the incidence, prevalence, or fatality rates of hereditary blood diseases (Khattak, & Saleem, 1992). βT is the most frequent genetically transmitted blood condition in Pakistan, affecting practically the whole country (Tasleem et al. 2007). While official data is unavailable in Pakistan, estimates of the number of babies with βT each year range from 5000 to 9000. Research indicates that the average carrier rate for βT in the general population is somewhat more than 5%. (Khattak, & Saleem, 1992) (Abolghasemi et al. 2007).

The increased frequency of βT cases can be attributed to inadequate education efforts and a lack of awareness (Asif, & Hassan, 2014). Many mothers with thalassemia trait are not aware that they are carriers, which can lead to the birth of a child with thalassemia major (Cao, & Kan, 2013). Given the autosomal recessive inheritance pattern, pre-marital carrier screening can be beneficial in prevention, and mandated programs have been implemented in numerous afflicted countries (Cousens

et al. 2010). In many countries it is being mandatory to analyzed the couple to screen and detect them from mutation causative age to take preventive measures as a part of patient therapy (Maheen et al. 2015). Globally many nations have initiate educational and diagnostic campaigns to prevent illness, while in Pakistan only two provinces have passed bills to provide financial support to national program that should be implemented to evaluate awareness and diagnostic screenings (Mutar et al. 2019). Therefore the aim of the study was to evaluate the level of awareness of the patient parents diagnosed with beta thalassemia.

METHODOLOGY:

Study Design:

This was a descriptive cross-sectional study conducted in different thalassemia centers in Swat. KPK The study was conducted in two thalassemia centers in Swat, which provide blood transfusion care facilities and services to all patients diagnosed with KPK at these centers. The study design was descriptive cross-sectional and was completed from February to March 2024. The study population consisted of parents of patients diagnosed with beta-thalassemia who received regular blood transfusions under the study conditions. Assuming that the total number of patients who were registered in both centers was the population, then using a 95% confidence level, 5% margin of error and 80% prevalence, the final sample size was 235 using a purposive sampling technique.

Data collection procedure:

The questionnaire was translated into Urdu (a local language) for better understanding of the questions, which were validated by subject specialists. I asked them and ticked the option. we stayed with the patient until all questions were answered, and we gave at least 20 minutes to each patient for a questionnaire.

Inclusion and exclusion criteria:

The inclusion criteria for the patient were: patient having a blood transfusion and diagnosed with beta-thalassemia after one year. Parents with cultural restrictions on communicating with the primary investigator or who are not willing to be voluntary participants are excluded from the study.

Data collection procedure:

The data collection process was started after getting permission from the administration of the study setting. Patients who were admitted for blood transfusions were approached and evaluated for inclusion criteria. The data was collected in two steps: In the first step, we collect the demographic data of the participants, which includes age, gender, and living status. In the second step, we collected data regarding awareness of informed consent. A valid and reliable questionnaire was adopted for data collection. The checklist contains 16 items with a dichotomous response (agree, neutral/disagree) (Muhammad et al. 2022).

Microsoft Excel and SPSS version 22 were used for the data analysis, which included descriptive statistics. For continuous variables, mean and standard deviation were computed; for categorical variables, frequency and percentages were used. Pearson correlation test was used to identify the association of awareness with demographic data.

The study was approved by the ethical review committee of Tasleem College of Nursing and Health Sciences, while permission was granted from the study setting for data collection, and informed consent was obtained from each participant to protect their ethical rights.

RESULTS

Socioeconomic status of the participants

In the current study majority of the parents were female 146 (62.1%), age group 31-40 years 164 (69.8%), house wife's 67 (28.5%), having monthly income 21000 to 50000 107 (45.5%), and parents with no education and matric 60 (25.5%), 60 (25.5%) respectively. (See table 1).

Table 1: Demographic data of the participants

	Frequency (235)	%
Gender		
Male	89	37.9
Female	146	62.1
Age		
21 - 30 years	29	12.3
31 - 40 years	164	69.8
41 - 50 years	34	14.5
51 and above	8	3.4
Occupation		
shopkeeper	46	19.6
Farmer	14	6.0
worker	72	30.6
Teacher	20	8.5
House wife	67	28.5
Driver	4	1.7
Overseas	4	1.7
Businessman	8	3.4
Monthly income		
20k and below	88	37.4
21 to 50k	107	45.5
51 to 100k	40	17.0
Education		
No education	60	25.5
Middle	37	15.7
Matric	60	25.5
intermediate	43	18.3
Bachelor	27	11.5
Masters and above	8	3.4

Level of awareness regarding thalassemia among parents of study participant

Table 2 illustrates that majority of the parents were aware regarding prohibited foods 124 (52%), first sign of

thalassemia 153 (65.1%), tests for prevention 104 (44.3%), complications regarding thalassemia 90 (38.3%), drugs not be taken by thalassemia patients 101 (43.0%), Regular lab investigation for the patient 128 (54.5%), thalassemia is genetic disorder 118 (50.2%), marriages in blood relations increase risk of Thalassemia 136 (57.9%), and HIV is blood borne diseases 88 (37.4%). (See table 2).


Table 2: Level of awareness among thalassemia patients parents

#	Questions	Correct	Don't, know	Incorrect
1	Know prohibited foods for thalassemia patient.	124 (52%)	69 (29.4%)	42 (17.9%)
2	Yellowish discoloration of skin is first sign of Thalassemia patient.	153 (65.1%)	24 (10.2%)	58 (24.7%)
3	For prevention method know the genetic investigation.	104 (44.3%)	66 (28.1%)	65 (27.7%)
4	Patients may experience splenomegaly as an outcome of their illness.	90 (38.3%)	65 (27.7%)	80 (34.0%)
5	Are you aware of bone marrow transplantation procedure?	84 (35.7%)	88 (37.4%)	63 (26.8%)
6	Transmission of Thalassemia is not occur through contact and food.	77 (32.8%)	53 (22.6%)	105 (44.7%)
7	Are there any drugs that the Thalassemia patients should not take?	101 (43.0%)	77 (32.8%)	57 (24.3%)
8	Through blood diabetes is not transmitted.	67 (28.5%)	53 (22.6%)	115 (48.9%)
9	Hemoglobin, liver and kidney function test is required regularly for their patients.	128 (54.5%)	42 (17.9%)	65 (27.7%)
10	You are considered as a carrier for Thalassemia	79 (33.6%)	88 (37.4%)	68 (28.9%)
11	Hepatitis can be transmitted via blood transfusion	72 (30.6%)	72 (30.6%)	91 (38.7%)
12	Hepatomegaly occurs as a outcome of their disease	86 (36.6%)	42 (17.9%)	107 (45.5%)
13	Is Thalassemia a genetic disease?	118 (50.2%)	53 (22.6%)	64 (27.2%)
14	Within the family marriage rises the likelihood of having an affected child	136 (57.9%)	50 (21.3%)	49 (20.9%)
15	HIV can be transmitted by blood transfusion	88 (37.4%)	85 (36.2%)	62 (26.4%)
16	If both of the carrier parents become pregnant, the chance of having an afflicted kid is 25%.	63 (26.8%)	83 (35.3%)	89 (37.9%)

Overall awareness of the participants

Figure 1 illustrates that majority of the patient level of awareness was good (42%), followed by poor awareness (31%), while 27% of the participant answer were neutral.

Figure 1: Overall level of awareness



Relationship of parents awareness with demographic variables

Table 3 illustrates that Parents awareness is significantly mild positive associated with gender, occupation and education, while negative mild associated with income while moderate negative with age. (See table 3).

Table 3: Association of awareness with demographic variables.

	1	2	3	4	5	6
1. Awareness	-	-.210**	.199**	.023	-.002	.185**
2. Age		-	-.109	-.121	.017	-.302**
3. Occupation			-	.163*	.276**	.185**
4. Income				-	.395**	-.033
5. Education					-	.156*
6. Awareness						-

**. Correlation is significant at the 0.01 level (2-tailed).

*. Correlation is significant at the 0.05 level (2-tailed).

DISCUSSION

The current study was conducted with the aim to collect the insight information of parents of Beta thalassemia patients regarding the diseases in Khyber pukhtankhwa.

In the present study majority of the parents were female 146 (62.1%), age group 31-40 years 164 (69.8%), house wife's 67 (28.5%), having monthly income 21000 to 50000 107 (45.5%), and parents with no education and matric 60 (25.5%), 60 (25.5%) respectively. Mothers have more access to the departmental resources as most of the attendant during transfusion with patient are mothers and uneducated mothers are due to cultural restrains in past.

A similar study conducted in Khyber pukhtankhwa Pakistan shows that the total participant were 220, where 252 (74.3%) were mothers, while 44 (25.7%). Higher number of parents were having poor level of knowledge regarding screening 237 (80.1%), while 265 (89.5%)

were have no education level (Shahzad et al. 2017).

In study conducted Pakistan shows that of the 88 patients that visited the Thalassemia Center, 76.5 percent were accompanied by their mothers, while the remaining patients were accompanied by their fathers. Out of 230 parents (both mothers and fathers), 74 (32.1%) were illiterate, 60 had 1-8 years of schooling (26%), 50 had 9-10 years of schooling (21.7%), 25 had 11-12 years of schooling, 14 (6.0%) had 13-14 years (10.7%) and only 7 had more than 15 years of schooling (3%). Of the 115 families surveyed, 39.1% earned less than PKR 6,000, 47% earned between PKR 6,000 and 15,000, and only 13.9% earned more than PKR 15,000 every month (Ansari et al. 2012). A study conducted in Lahore Pakistan reveals that among the total 410 families, 300 (73.2%) coming from public and 110 (26.8%) from private centers. There were 179 (43.7%) fathers and 231 (56.3%) moms who attended with their children, ranging in age from 20 to 60 years. There were 279 rural families (69%) and 127 urban families (31%). In terms of parental education, 18.3% (75 out of 179) fathers and 32.9% (135 out of 231) women did not receive a formal education (ISHFAQ et al. 2016).

In the current study 118 (50.2%) of the participant were aware that thalassemia is genetic disorder, it may be due to minor cases in towns that concern the family member to know about the diseases and also when the child is diagnosed first time. Moreover the majority of the parents were aware regarding prohibited foods 124 (52%), first sign of thalassemia 153 (65.1%), tests for prevention 104 (44.3%), complications regarding thalassemia 90 (38.3%), drugs not to be taken by thalassemia patients 101 (43.0%), regular lab investigation for the patient 128 (54.5%), marriages in blood relations increase the risk of thalassemia 136 (57.9%), and HIV is a blood-borne disease. A study completed in Multan Pakistan support our findings were 60(50.0%) of the participants were aware that it is inherited diseases, approximately 29.2% (n=35) of the parents believed that a thalassemia patient may survive without therapy, while most parents did not know about it. 64.2% and 6.7% of parents answered that they did not know. Similarly, 89.2% of parents know that this disease can be identified through blood tests, and a majority of 91.7% expect that blood transfusion is a cure for this disease, while 49.2% respondents believed that marriage is an important risk factor for thalassemia in their children (Saleem et al. 2021). Another Pakistani study reveals that Out of the accompanying parents, 52 (44.6%) were aware that thalassemia is an inherited blood illness. However, 63 (55.6%) were unaware of its etiology. Only 37 (32.2%) were aware that both parents could be carriers for the condition in their children (Ansari et al. 2012). A study conducted in Iraq shows different findings from our study where 54% of the parents were not aware that it is hereditary diseases, moreover majority 94.8% of the parents were aware regarding type of food should not be taken, 86% about the sign of thalassemic patient, 83.7% about genetic

tests, 80.1% about bone marrow transplantation, 78.2% about transmission of diseases, 77.7% about drugs not to be used by the patient (Muhammad et al. 2022).

In the current study majority of the participant level of awareness was good (42%), followed by poor awareness (31%), while 27% of the participant answer were neutral. It may be due to regular transfusion session attended in blood transfusion centers, interaction with doctors and being a special patient in the community. A Pakistani study found that 82% of parents were aware of their child's thalassemia diagnosis and 80% knew it was a blood illness (Ghafoor et al. 2016). A study conducted in Islamabad Pakistan reports that the majority of the parents, 274 (66.8%), were unaware of thalassaemia prior to their first affected kid. Similarly, 333 (81.2%) participants held the incorrect belief that the disease was genetic (ISHFAQ et al. 2016). A study conducted in Khyber pukhtankhwa shows that only 16 out of 162 patients with family history of β T major were aware of prenatal screening. As a result, many families with β T major are unaware of their condition (Shahzad et al. 2017).

In the current study, parental awareness is significantly moderate positive related to gender, occupation and education, while negative moderate related to income, while moderate negative related to age. Ghafoor et al. found that 60% of parents knew nothing about thalassemia and 69% were uninformed (Arif et al. 2008). The majority of women (89.5%) were uneducated, which explains the lack of information on prenatal screening for β T major. Our survey found that the literacy rate is higher than in Karachi (57%) (Ishfaq et al. 2015).

There were certain limitation in form of sampling technique, study design and limited to a province that could be generalizable to all the country.

CONCLUSION

The study concluded that majority of the parents level of awareness is good regarding common factors that are associated with β T. Parents' understanding regarding thalassemia carrier screening and prenatal diagnosis was inadequate. To address the issue, it's crucial to start a national level education campaign for the public to raise the awareness among families with thalassemic conditions.

Conflict of interest: The authors declare that there is no conflict of interest.

REFERENCES

- Abolghasemi, H., Amid, A., Zeinali, S., Radfar, M. H., Eshghi, P., Rahiminejad, M. S., ... & Hoorfar, H. (2007). Thalassemia in Iran: epidemiology, prevention, and management. *Journal of Pediatric Hematology/Oncology*, 29(4), 233-238.
- Ansari, S. H., Shamsi, T. S., Ashraf, M., Farzana, T., Bohray, M., Perveen, K., ... & Raza, F. (2012). Molecular epidemiology of β -thalassemia in Pakistan: Far reaching implications. *Indian journal of human genetics*, 18(2), 193.
- Arif, F., Fayyaz, J., & Hamid, A. (2008). Awareness among parents of children with thalassemia major. *J Pak Med Assoc*, 58(11), 621-624.
- Asif, N., & Hassan, K. (2014). Prevention of beta thalassemia in Pakistan. *J Islam Med Dent Coll*, 3(2), 46-7.
- Cao, A., & Kan, Y. W. (2013). The prevention of thalassemia. *Cold Spring Harbor perspectives in medicine*, 3(2), a011775.
- Cousens, N. E., Gaff, C. L., Metcalfe, S. A., & Delatycki, M. B. (2010). Carrier screening for beta-thalassaemia: a review of international practice. *European journal of human genetics*, 18(10), 1077-1083.
- Cousens, N. E., Gaff, C. L., Metcalfe, S. A., & Delatycki, M. B. (2010). Carrier screening for beta-thalassaemia: a review of international practice. *European journal of human genetics*, 18(10), 1077-1083.
- Ghafoor, M. B., Leghari, M. S., Mustafa, G., & Naveed, S. (2016). Level of awareness about thalassemia among parents of thalassaemic children. *Journal of Rawalpindi Medical College*, 20(3).
- Hay, S. I., Jayaraman, S. P., Truelsen, T., Sorensen, R. J., Millea, A., Giussani, G., & Beghi, E. (2017). GBD 2015 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015 (vol 388, pg 1545, 2016).
- Ishaq, F., Hasnain Abid, F. K., Akhtar, A., & Mahmood, S. (2012). Awareness among parents of $\beta\beta$ -Thalassemia major patients, regarding prenatal diagnosis and premarital screening. *Journal of the College of Physicians and Surgeons Pakistan*, 22(4), 218-21.
- ISHFAQ, K., AHMAD, T., NAEEM, S. B., ALI, J., & ZAINAB, S. (2016). THE KNOWLEDGE OF PARENTS HAVING THALASSEMIA CHILD. *Isra Medical Journal*, 8(2).
- Ishfaq, K., Hashmi, M., & Naeem, S. B. (2015). Mothers' awareness and experiences of having a thalassemic child: a qualitative approach. *Pakistan Journal of Applied Social Sciences*, 2(1), 35-53.
- Khattak, M. F., & Saleem, M. (1992). Prevalence of heterozygous beta-thalassemia in northern areas of Pakistan. *JPMA. The Journal of the Pakistan Medical Association*, 42(2), 32-34.
- Maheen, H., Malik, F., Siddique, B., & Qidwai, A. (2015). Assessing parental knowledge about thalassemia in a thalassemia center of Karachi, Pakistan. *Journal of genetic counseling*, 24(6), 945-951.
- Muhammad, L., Wajid, K. K., Afridi, I., Ullah, S., Khan, A., Muhammad, A., ... & Muhammad II, A. (2022). Awareness Among Educated and Uneducated Parents of Beta-Thalassemia Major Patients About Antenatal Screening. *Cureus*, 14(8).
- Mutar, M. T., Majid, M., Jaleel, A., Saad, A., Abdulmorteza, A., & Talib, H. (2019). Awareness among parents of beta thalassemia major and intermedia patients in three centers in Baghdad and Al-Nasiriyah, Iraq in 2017. *International Journal of Medical Students*, 7(1), 6-10.
- Saleem, N., Anwar, A., Shahid, N. U. A., Saleem, R., Saleem, Z., Asghar, H., ... & Iqbal, S. (2021). Perception of parents of thalassemic child to thalassemia in Pakistan. *Cureus*, 13(8).
- Shahzad, A., Rafiq, N., Ullah, I., Asad, M. J., Ahmad, M. S., & Waheed, U. (2017). Knowledge, attitude and practices (KAP) of the families of β -thalassaemia children in thalassaemia centers of Rawalpindi and Islamabad, Pakistan. *Blood transfusion (BT)*, 92, 22-4.
- Tasleem, S., Tasleem, H., Siddiqui, M. A., Adil, M. M., & Rashid, Y. (2007). Prenatal diagnosis of β -thalassaemia by chorionic villous sampling. *JPMA*, 57(11), 528-31.
- Thalassemias 2012. National Heart, Lung, and Blood Institute: Thalassemias. 2012.
- Uthman, O. A. (2016). Global, regional, and national life expectancy, all-cause and cause-specific mortality for 249 causes of death, 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*, 388(10053), 1459-1544.