

Knowledge, Attitude and Practice on Dental Caries and Oral Hygiene among Medical Students Islamabad Pakistan

Muhammad Asif¹, Ume Habiba¹, Naila Iftikhar³, Muhammad Ans², Sadia Sana³, Benish Ali⁴, Hina Anwar⁵

Abstract: The objective of the study was to evaluate the awareness, attitudes and practices on oral hygiene and dental caries among medical students studying at Rawal Medical and Dental College, Islamabad. It was a descriptive cross-sectional study carried out at Rawal Medical and Dental College, Islamabad in a time period of four months i.e. from 1st January 2018 to 30th April 2018 by using convenient sampling technique. 350 students of MBBS (1st to 5th year) of Rawal Medical and Dental College were enrolled in this research study. A total fourteen-item close-ended questionnaire, consisting of 4 items for knowledge about dental caries (including sugar, bacteria, not brushing causes caries, benefits of fluoride on dental health), 5 for each attitudes and practice (including visit on dental clinic for dental checkup, and frequency of brushing, tooth paste type, brushing time, brushing methods and materials). The results of this study revealed that out of 330 medical students, 185 were female and 145 were male. Male students had significantly greater awareness about dental caries ($p < 0.05$). Majority of these medical students (46%) had tendency of visiting once in year for dental checkup. The maximum number of medical students had once per day (60%) brushing routine in early morning (54.2%) with tooth paste (52%) containing fluoride content. The knowledge on dental caries was found as significantly associated with the use of fluoridated dentrifice, brushing practices, brushing methods and brushing materials ($p < 0.05$). The current study concluded that awareness and practice status of dental caries and oral hygiene in students of medical profession was found to be satisfactory. However further oral health education programs must be carried out for the control and prevention of oral ailments and better dental hygiene among the dental students.

Keywords: Oral hygiene, Dental caries, Medical student, Attitude, Knowledge

- 1 Department of public health, Directorate of Medical Sciences, Government College University, Faisalabad
- 2 College of Physical therapy, Directorate of Medical Sciences, Government College University, Faisalabad
- 3 College of Allied Health professionals, Department of public health, Directorate of Medical Sciences, Government College University, Faisalabad
- 4 Department of Orthotics and prosthetics, Department of public health, Directorate of Medical Sciences, Government College University, Faisalabad
- 5 Department of Eastern Medicine, Directorate of Medical Sciences, Government College University, Faisalabad

Corresponding author's email:
asif.msph094@gmail.com

INTRODUCTION

Oral health is taken as fundamental to general health and well-being. The knowledge related to oral health is understood to be a necessary prerequisite for health related behavior (1). Oral hygiene focus on the practices that retain the mouth healthy and hygienic by brushing and flossing to prevent tooth decay and gum disease (2). A healthy mouth facilitates an person to talk, socialize and eat without facing any active disease, embarrassment and distress (3, 4). Several oral disorders have significant effects on status of general health, whereas systemic disorders may put a common effect on oral health and are connected with significant pain, anxiety and impaired social performance (5, 6). Worldwide, dental caries is considered as a key public health problem due to its high prevalence and considerable social impact in essential life activities. The microbial community of caries is varied and contains several facultative and obligate anaerobic bacteria. Streptococcus mutans is considered to be the main etiological agent of dental caries (7). There are three factors responsible for causing dental caries; diet, bacteria and host (8). Many studies have been carried out to detect risk indicators of dental

caries to find out high-risk individuals (9). Pakistani people are less aware of oral health due to socio-economic issues and insufficient health care resources (10). In Pakistan, the oral health system is currently in transition phase and also needs significant resources and facilities. The lack of dental awareness has a significant influence on two most prevalent dental diseases; periodontal diseases and dental caries (11).

The highly important time period of a student's life is spent at school and college which is the place that develops their attitudes, habits and beliefs for lifetime (12). Organization can play a leading role in conveying awareness of the causes and avoidance of most prevalent oral disorders which delivers an ideal situation for encouraging good oral health activities among all age groups. The principle of risk-based management of caries in clinical education has been recommended and is part of the undergraduate curriculum in recent years (13). There are very limited studies which have collected data on knowledge of dental caries among children and health professionals but the information is scanty with concern to medical students.

Due to limitation of studies on attitude and behaviour about oral health among University medical students, the current study is vital in health services. In scientific literature there is deficiency of data on knowledge, attitude and behaviour about dental health among medical students. The current research study aims to evaluate the level of awareness, attitudes and practices on dental caries and oral hygiene among medical students studying at Rawal Medical and Dental College, Islamabad.

METHODOLOGY

It was a descriptive, cross-sectional study carried at Rawal Medical and Dental College, within a period of four months i.e. from 1st January 2018 to 30th April 2018. By using convenient sampling technique, a self-administered standard structured questionnaire was distributed among 350 medical students studying in Bachelor of Medicine and Bachelor of Surgery (MBBS) programme to assess the knowledge of oral hygiene and dental caries, out of which 330 students of medical profession completed the questionnaire.

A total fourteen-item close-ended questionnaire, consisting of 4 items for knowledge about dental caries (including sugar causes caries, bacteria causes caries, not brushing causes caries, benefits of fluoride on dental health), 5 for each attitudes and practice (including visit on dental clinic for dental check up, once in three months, only if in pain, once in years and never visited and frequency of brushing, tooth paste type, brushing time, brushing methods and materials). The questionnaire was constructed in English language.

Only the students of basic science and clinical science

studying MBBS programme were included, whereas health professionals such as consultant, medical officer, paramedics and nursing professionals were excluded. Verbal informed consent was taken and the objective of the research study was elaborated to the participants evidently with assurance of confidentiality.

STATISTICAL ANALYSIS

The data were explored with the help of SPSS version 16.0 statistical software. Frequency distribution was used to describe the distribution of knowledge on dental caries cases in the study sample. The Chi-square test was utilized to check for the relation between knowledge on dental caries with gender, use of fluoridated dentrifice, brushing habit, brushing method and brushing material. The p-value was calculated as <0.05 showing statistically significant results.

RESULTS

Out of 330 medical students, 185 (56.1%) were female and 145 (43.9%) were male. Among all respondents, male respondents had more awareness than female respondents about dental caries. 78 (23.63%), 83 (25.25%) and 94 (28.48%) male respondents had knowledge about sugar, not brushing and bacteria respectively that causes caries and 58 (17.57%) had knowledge about benefits of fluoride on dental health and rest of them did not know about the subject matter. Similarly, 25 (7.57%), 33 (10%) and 24 (7.27%) female respondents were having knowledge about sugar, not brushing and bacteria respectively that causes caries and 13 (3.93%) had knowledge about benefits of fluoride on dental health and rest of them did not know about the subject matter. The results are shown in table I.

Table I. Knowledge regarding dental caries among the study population (n = 330)

Causes	Knowledge on dental caries			
	Yes		No	
	Male	Female	Male	Female
	No. (%)	No. (%)	No. (%)	No. (%)
Sugar	78 (23.63)	25 (7.57)	67 (20.30)	160 (48.48)
Not brushing	83 (25.25)	33 (10)	62 (18.78)	152 (46.06)
Bacteria	94 (28.48)	24 (7.27)	52 (15.75)	160 (48.48)
Benefits of fluoride on dental health	58 (17.57)	13 (3.93)	87 (26.36)	171 (51.81)

Out of total population, 123 (37.3%) respondents had knowledge on dental caries and among them 95 (65.5%) were male and 28 (15.1%) were female respondents. The

knowledge on dental caries was found higher in males than females. At 5% significance level the result was statistically significant (Table II).

Out of 330 respondents, 173 (52%) respondents used fluoridated dentrifice and 157 (48%) respondents not used fluoridated dentrifice.

Among the total participants, 79 (45.7%) participants had awareness about dental caries who used fluoridated

dentrifice and those who did not use it had 44 (28%) awareness about dental caries. The awareness about dental caries was noted to be maximum among the respondents who utilized fluoridated dentrifice. At 5% significance level the result was statistically significant (Table II).

Table II. Association between gender, use of fluoridated dentrifice and knowledge of dental caries (n = 330)

	Knowledge on dental caries			p-value
	Yes	No	Total	
	No. (%)	No. (%)	(No)	
Gender				
Male	95 (65.5)	50 (34.5)	145	0.0001
Female	28 (15.1)	157 (84.8)	185	
Total	123 (37.3)	207 (62.7)	330	
Fluoridated dentrifice				
Used	79 (45.7)	94 (54.3)	173	0.001
Not used	44 (28.0)	113 (72.0)	157	
Total	123 (37.3)	207 (62.7)	330	

Among all respondents, 265 (39%) respondents visited dental clinic for dental checkup, 38 (6%) were once in three months, 49 (7%) were only if in pain, 309 (46%) once in year and 12 (2%) were never visited.

Out of total respondents, 11 (3.3%), 198 (60%) and 121 (36.7%) respondents were found as brushing their tooth occasionally, once per day and twice per day respectively. Similarly, 50 (15.2%), 101 (30.6%) and 179 (54.2%) respondents were brushing their tooth before and after eating, after eating and in early morning respectively.

Among all, most of the respondents had brushing habit once per day and brushed their tooth in early morning.

Among the respondents, 36.4%, 37.9% and 36.4% respondents had knowledge on dental caries with occasional, once per day and twice per day brushing habit respectively. The awareness about dental caries was noted to be maximum in those respondents who had brushing habit once per day. The result was not statistically significant at 5% level of significance. Similarly, 58%, 28.7% and 36.3% of participants had awareness about dental caries with before and after eating, after eating and early morning brushing time respectively. The awareness about dental caries was found to be maximum among the respondents who had before and after eating brushing habit. The result was statistically significant at 5% level of significance (Table III).

Table III. Association between brushing habit and knowledge of dental caries (n = 330)

Brushing habit	Knowledge on dental caries			p-value
	Yes	No	Total	
	No. (%)	No. (%)	(No)	
Occasional	4 (36.4)	7 (63.6)	11	0.962
Once per day	75 (37.9)	123 (62.1)	198	
Twice per day	44 (36.4)	77 (63.6)	121	
Before and after eating	29 (58)	21 (42.0)	50	0.002
After eating	29 (28.7)	72 (71.3)	101	
Early morning	65 (36.3)	114 (63.7)	179	

Among all respondents, 265 (39%) respondents visited dental clinic for dental check up, 38 (6%) were once in three months, 49 (7%) were only if in pain, 309 (46%) once in year and 12 (2%) were never visited. Out of total respondents, 11 (3.3%), 198 (60%) and 121 (36.7%) respondents were found as brushing their tooth occasionally, once per day and twice per day respectively. Similarly, 50 (15.2%), 101 (30.6%) and 179 (54.2%) respondents were brushing their tooth before and after eating, after eating and in early morning respectively.

Among all, most of the respondents had brushing habit once per day and brushed their tooth in early morning.

Among the respondents, 36.4%, 37.9% and 36.4% respondents had knowledge on dental caries with occasional, once per day and twice per day brushing habit respectively. The awareness about dental caries was noted to be maximum in those respondents who had brushing habit once per day. The result was not statistically significant at 5% level of significance. Similarly, 58%, 28.7% and 36.3% of participants had awareness about dental caries with before and after

eating, after eating and early morning brushing time respectively. The awareness about dental caries was found to be maximum among the respondents who had before and after eating brushing habit. The result was statistically significant at 5% level of significance (Table III).

Among all respondents brushing method, 108 (33%) respondents followed horizontal, 64 (19%) vertical, 158 (48%) mixed methods during brushing. Similarly, 66 (20%) respondents used brush and tooth powder, 166 (50.3%) used brush and tooth paste, 52 (15.8%) used finger and tooth powder and 46 (13.9%) used dattiwan and others material for brushing.

The awareness on dental caries was revealed to be maximum among those respondents who followed vertical brushing methods and who used dattiwan and others materials during brushing. At 5% significance level the result was statistically significant (Table IV).

DISCUSSION

Oral health is a basic element of the general health of a person and has been declared a vital problem

Table IV. Association between knowledge on dental caries and brushing practices (n = 330)

Brushing habit	Knowledge on dental caries			p-value
	Yes	No	Total	
	No. (%)	No. (%)	(No)	
Horizontal	26 (24.1)	82 (75.9)	108	0.002
Vertical	31 (48.4)	123 (62.1)	64	
Mixed	66 (41.8)	92 (58.2)	158	
Brushing materials				
Brush and Tooth Powder	29 (58)	21 (42.0)	50	0.0001
Brush and Tooth Paste	29 (28.7)	72 (71.3)	101	
Fingerand Tooth Powder	65 (36.3)	114 (63.7)	179	
Dattiwanand Others	34 (73.9)	12 (26.1)	46	

of public health with a significant social impact (14). As William Osler said mouth is the mirror of general health of human beings (15). Optimal health cannot be accomplished independent of oral health. Unhealthy oral conditions may very badly disturb general health of the population and some medical disorders may have an undesirable influence on oral health (16).

Oral diseases are more prevalent in the developing nations. In Pakistan, there is limited data available on the oral health status of medical students. Therefore, the present research study was focused on the dental caries and knowledge regarding oral hygiene among the student of medical profession. Knowledge acquisition regarding oral health includes complex cognitive processes: reasoning, association, learning, communication and perception of the conditions about oral health (17). It was found that the total knowledge on dental caries was higher in male than female which was found statistically significant at 5% level of significance ($p=0.0001$). This may be due to knowledge on oral hygiene in male than female. The results of a Pakistani study conducted by Mehmood et al revealed that the knowledge level of 95% of students was satisfactory. Among them 47% were male and 48% were female students (18).

This study highlights that 31.2% respondents knew that sugar causes caries, 35.2% knew that not brushing teeth causes caries, 35.8% knew that bacteria cause caries and 21.6% had knowledge about benefits of fluoride on dental health. The knowledge on dental caries in those who used fluoridated dentrifice was 45.7% which was

found statistically significant at 5% level of significance ($p=0.001$). Fluoride has an anti-cariogenic effect that prevents caries and can even reverse the initial phases of teeth decay (19).

The present research study showed the highest number of respondents had habit of visiting once in a year for dental check-up which indicates the negative attitude towards dental care. The results of a study conducted by Baseer et al revealed that the attitude toward visiting the dentist varied among different health professionals like 50% of technicians and 52.7% of nurses reported that they regularly go to the dentist for dental checkup. In the same way it was found that 66.7% of the medical students go to consult the dentist whenever they suffer from tooth ache. 54.5% doctors and 45.8% pharmacists visited the dentist occasionally (20).

This study found that the maximum number of respondents had brushing habit once per day and also had more knowledge on caries, which was not statistically significant at 5% level of significance ($p=0.962$). This may be due to lack of knowledge about brushing twice a day is good for maintaining the proper oral hygiene. A similar finding was also obtained in study conducted by Baral et al [21]. In a Pakistani study conducted by Mehmood et al, it was found that 41.7% of students use to clean their teeth once, 32.2% clean twice, 17.7% students clean their teeth three times and 8% clean their teeth more than three times a day (18).

This present study also establishes the awareness about dental caries was maximum among the respondents who

had before and after eating brushing habit which was statistically significant at 5% level of significance ($p=0.002$). In this study, the highest number of respondents followed mixed brushing methods. A study conducted by Baral et al also obtained the similar findings (21). The present study also found the knowledge and awareness about dental caries was found maximum among those respondents who followed vertical brushing methods and who used dattiwan and others materials during brushing, which is statistically significant at 5% level of significance. The vertical brushing method considers manual dexterity to rotate bristles from the gums towards and across the teeth which helps to prevent from caries. Dattiwan is a stem or root of the plant which becomes brushlike after treatment and is suitable for oral hygiene (22). Although the medical students have good knowledge of benefits of dattiwan using, but the practice of dattiwan was not so frequent among the students of medical college.

The current research study data showed the knowledge, awareness, attitude and habit practice status of medical profession students which helps to achieve optimal oral health care. Further oral health educational programs in

the schools, colleges, Universities and community must be arranged on regular basis for the control and prevention of oral diseases.

CONCLUSION

This study was the formal assessment of dental health knowledge and related behavior of medical students. The present study concluded that male respondents showed relatively good knowledge of oral hygiene and dental caries than female respondents which was found to be statistically significant. It is possible that female had not particular knowledge about dietary factors causing caries. However carrying knowledge and awareness does not assure that it will be efficiently utilized; this requires effective campaigning within community through various medical associations and organizations. Adequate knowledge, good behaviors and oral hygiene practices among medical students are further suggested to be enhanced with positive attitudes, comprehensive knowledge, and healthy behavior in relation to the oral health.

REFERENCES

1. Al-Ansari J, Honkata E, Honkata S. Oral health knowledge and behavior among male health sciences college students in Kuwait. *BMC Oral Health*. 2013; 3(12):2-5.
2. Emmanuel A, Changendo E. Oral health related behaviour, knowledge, attitudes and beliefs among secondary school students in Iringa municipality. *DMSJ*. 2010; 17(1): 24-30.
3. Australian Research Centre for Population Oral Health, the University of Adelaide, South Australia. Periodontal diseases in the Australian adult population. *Aust Dent J*. 2009; 54(3):390-393.
4. Zhu L, Petersen PE, Wang HY, Bian JY, Zhang BX. Oral health knowledge, attitudes and behaviour of adults in China. *Int Dent J*. 2015; 55(3):231-241.
5. Kelly M, Steele J, Nuttall N, Bradnock G, Morris J, Nunn J, et al. Adult dental health survey: oral health in the United Kingdom 2000. London. 45-50.
6. Chen MS, Andersen RM, Barmes DE, et al: Comparing oral health care systems: a second international collaborative study. Geneva: World Health Organization; 2005. 35-41
7. Yadav K, Prakash S. Antibigram profiles against polymicrobial pathogens among dental caries patients at Janaki Medical College teaching hospital, Nepal. *International Journal of Applied Dental Sciences*. 2015; 1(4): 156-162.
8. Kisumbi BK, Kaimenyi JT, Wakiaga JM. Knowledge on treatment modalities and attitude of Nairobi University students towards dental care. *Indian Journal of Dental Research*. 2015; 6(4):133-36.
9. American Dental Association Council on Scientific Affairs. Professionally applied topical fluoride: Evidence-based clinical recommendations. *Journal of American Dental Association*. 2012; 137(9): 1151-1159.
10. Yee R, Mishra R. Nepal National Oral Health Pathfinder Survey 2004. Kathmandu, Nepal: Oral Health Focal Point; Ministry of Health; HMG Nepal. 2004. 105-111
11. Zhang J, Xuan D, Fan W, Zhang X, Dibart S, De Vizio W, Panagakos F, Zhang YP. Severity and prevalence of plaque-induced gingivitis in the Chinese population. *Compend Contin Educ Dent*. 2010; 31(8):624-629.
12. Dawani N, Afaqand A, Bilal S. Oral health knowledge, attitude and practices amongst teachers of public school set-up of Karachi, Pakistan. *J Dow Univ Health Sci Karachi*. 2013; 7(1): 15-19.
13. Dodds MW, Suddick RP. Caries risk assessment for determination of focus and intensity of prevention in a dental school clinic. *Journal of Dental Education*. 1995; 59(8): 945-956.
14. The World Oral Health Report .Continuous improvement in the oral health in the 21st century- the approach of the WHO Global Oral Health Programme. 2003. 67-72
15. Sumi Y, Nakamura Y, Nagosa, et al. Attitudes to oral care among caregivers in Japanese nursing homes. *Gerodontology*. 2001; 18(1): 1-6.
16. Radha G, Shaik Hyder Ali KH, Pushpanjali K. Knowledge and attitude and practice of oral health among nursing staff and nursing students of Bangalore city. *Journal of Indian Association of Public Health Dentistry*. 2008; (11): 17-21.
17. Skaret E, Espelid I, Skeieb MS, & Haugejorden O. Parental Beliefs and Attitudes towards Child Caries Prevention: Assessing Consistency and Validity in a Longitudinal Design. *BMC Oral Health*. 2008; 8(1): 8-19.
18. Mehmood R, Rasheed S, Ijaz M. KNOWLEDGE, ATTITUDE AND PRACTICE REGARDING ORAL HEALTH AMONG SECONDARY SCHOOL STUDENTS

- OF AZAD KASHMIR, PAKISTAN. Pakistan Journal of Public Health 2018; 8(1): 32-36.
19. Whelton H. Overview of the impact of changing global patterns of dental caries experience on caries clinical trials. J Dent Res. 2014; 83 (1): 29-34.
20. Baseer MA, Alenazy MS, AlAsqah M, AlGabbani M, Mehkari A. Oral health knowledge, attitude and practices among health professionals in King Fahad Medical City, Riyadh. Dental Research Journal. 2012;9(4):386-392.
21. Wright FA. Children's perception of vulnerability to illness and dental disease. Community Dent Oral Epidemiol. 2011;10(1): 29-32.
22. Baral P, Bhattarai C, Poudel PP, Banstola D, Roy S, Hada S and Hiremath SS. A Study on Oral Hygiene Practice among School Children of Pokhara Municipality. J of GMC-Nepal. 2009; 2(1):37-38.